

# A review of the practice of surgical antibiotic prophylaxis

Otaigbe II<sup>1, 2</sup>

<sup>1</sup>Department of Medical Microbiology and Parasitology, School of Basic Clinical Sciences, Benjamin Carson (Snr) College of Health and Medical Sciences, Babcock University, Ilishan Remo, Ogun State, Nigeria.

<sup>2</sup>Department of Medical Microbiology and Parasitology, Babcock University Teaching Hospital, Ilishan Remo, Ogun State, Nigeria.

Submitted: 24<sup>th</sup> July 2023

Accepted: 6<sup>th</sup> October 2023

Published: 31<sup>st</sup> December 2023

: Orcid ID

## Abstract

**Background:** Surgical Antibiotic Prophylaxis (SAP) is defined as the use of antibiotics to prevent infections at a surgical site. The appropriate use of SAP is an effective and efficient modality of preventing surgical site infections.

**Main body:** The practice of SAP is however fraught with several challenges e.g., diverse opinions among surgeons regarding the duration of surgical prophylaxis, a dearth of institutional guidelines, and poor compliance with existing guidelines. Unfortunately, the inappropriate use of SAP results in adverse clinical (e.g., increased length of hospitalization, occurrence of surgical site infections, morbidity, etc.) and economic consequences (e.g., increased healthcare costs) in surgical patients.

**Conclusion:** This review article discusses the practice of SAP and the challenges regarding SAP. Finally, some recommendations to overcome the challenges are discussed.

**Keywords:** Surgical antibiotic prophylaxis, Surgical site infection, Guidelines, Compliance, Quality improvement

## Background

Surgical antibiotic prophylaxis (SAP) is defined as the use of antibiotics to prevent infections at a surgical site (1). It involves the initial administration of a short course of an antimicrobial agent before surgery to prevent surgical site infections (2). The appropriate use of SAP is an effective and efficient modality of preventing surgical site infections (SSI) (1, 3). Antibiotics however carry the risk of adverse effects and drug resistance (4). The choice of an antibiotic for SAP should therefore ensure coverage of organisms likely to cause an infection at the surgical site and be influenced by the strength of association between the antibiotic used and these adverse effects (5). The above

statement is best achieved with regular audits of the practice of SAP and the design of local antibiotic formularies which would ensure that the most appropriate antibiotic, dose, timing of administration, and duration are used for effective prophylaxis (5).

The use of SAP takes into consideration the following: the need to decide if the surgical procedure requires the use of antibiotic prophylaxis; a knowledge of the bacterial flora most likely to cause a surgical site infection; the choice of an antibiotic, based on the steps above, with the narrowest antibacterial spectrum required; the choice of the less expensive drug if two drugs are otherwise of the equal antibacterial spectrum,

Correspondence:

Otaigbe, Idemudia I

Department of Medical Microbiology,

School of Basic Clinical Sciences, Benjamin Carson (Snr) College of Health and Medical Sciences,

Babcock University/Babcock University Teaching Hospital,

Ilishan Remo, Ogun State, Nigeria

+2348024406763, [otaigbei@babcock.edu.ng](mailto:otaigbei@babcock.edu.ng)

efficacy, toxicity, and ease of administration; the administration of the antibiotic at the right time and the right dose; the use of surgical prophylaxis for a short period (one dose if surgery of four hours duration or less); the avoidance of antibiotics likely to be of use in the treatment of serious sepsis; the knowledge by the surgical team that SAP is not an attempt to overcome poor surgical technique and the presence of (and periodic reviews of) local guidelines on SAP (1).

To ensure optimum best practices in surgery and quality healthcare delivery several attempts have been made at institutional, national, and international levels to develop guidelines and recommendations for SAP (4). Several studies have however documented poor compliance with existing guidelines. Sadly, the adverse effects (e.g., the occurrence of surgical site infections, inappropriate use of antibiotics, emergence of drug resistance, increased costs to the patient, the risk of drug toxicity etc.) of poor guideline compliance are borne predominantly by surgical patients (6). For example, a study done by Fukatsu et al found a correlation between prolonged use of surgical prophylaxis and the development of methicillin-resistant *Staphylococcus aureus* (7). While some sections in this paper, discuss the epidemiology, risk factors and microbiology of surgical site infections the focus of this narrative review is the practice of SAP.

## Main body

### Methods

This paper is a narrative review of the practice of surgical antibiotic prophylaxis. It essentially involved a literature review of existing theories and recommendations regarding the practice of surgical antibiotic prophylaxis. In line with narrative reviews, there was no predetermined research question nor specified search strategy or protocol regarding the topic i.e. surgical antibiotic prophylaxis.

### *The concept of surgical antibiotic prophylaxis*

Surgical antibiotic prophylaxis (SAP) is defined as the use of antibiotics to prevent infections at a surgical site (1). It refers to a very brief course of an antimicrobial agent commenced just before the onset of an elective surgical procedure (1). It is not an attempt to sterilize tissues involved in the surgery nor does it seek to prevent surgical site infections caused by contamination, in the post-operative period (1). Rather it is a decisive and properly timed process which is indicated in an elective operation in which skin incisions are closed during surgery (1). It serves to reduce the microbial

burden of intraoperative contamination to a level that cannot overwhelm the patient's host defences (1). Its tacit aim is to prevent surgical site infections through the administration of safe, cost-effective antimicrobials which possess a spectrum of activity against pathogens that are likely to cause an infection at the surgical site (1).

The appropriate and timely administration of antibiotic prophylaxis has been shown to reduce the incidence of surgical site infections (1). Dettenkofer et al (2002) state that approximately 30 to 50% of antibiotic use in hospital practice is for surgical prophylaxis and that between 30% to 90% of this prophylaxis is inappropriate (e.g., inappropriate timing or duration) (8). Such injudicious use of antibiotics also increases the selective pressure favouring the emergence of antimicrobial resistance (1).

Ideally, an anti-infective drug for surgical prophylaxis should achieve the following goals: it should prevent the occurrence of a postoperative infection at the surgical site; prevent postoperative infectious morbidity and mortality; it should reduce the duration and cost of health care; it should produce no adverse effects, and have no adverse consequences for the microbial flora of the patient or the hospital (8, 9). To achieve these goals, an anti-infective drug should be: active against the pathogens most likely to contaminate the wound; given in an appropriate dosage and at a time that ensures adequate concentrations at the incision site during the period of potential contamination; safe; and administered for the shortest effective period to minimize adverse effects, development of resistance, and cost (8, 9).

### *Historical background*

In the nineteenth century, the works of Louis Pasteur and Joseph Lister laid the background for the introduction of asepsis and prophylaxis in surgical practice (4).

While Louis Pasteur discovered that microbes were responsible for disease and putrefaction Joseph Lister's discovery of aseptic procedures was instrumental in wound healing and reduced mortality rates associated with surgeries at this time (4).

Despite these major discoveries, the use of surgical prophylaxis was fraught with opposition from surgeons and academicians involved in the teaching and practice of surgery (4). For example, in 1880, nearly 15 years after the discoveries of Pasteur and Lister, a surgeon called William Halstead was ordered from the operating theatre for daring to challenge a senior surgeon's non-compliance with aseptic surgical techniques (4).

Furthermore, many surgeons could not reach a consensus on the need for prophylaxis in surgery (3). However, studies done between 1961 and 1963 by investigators in Cincinnati and Boston showed the isolation of *Staphylococcus aureus* from the operative field, despite the application of standard aseptic techniques. The investigators concluded that while standard aseptic techniques could decrease but not eliminate bacterial contamination of the surgical field, the addition of surgical prophylaxis (to standard aseptic techniques) is vital in preventing infections at the surgical site (10, 11, 12).

The second discovery showed that early administration of surgical prophylaxis was associated with the prevention of infection at the surgical site (13, 14, 15). The efficacy of prophylactic antibiotics has been verified and peri-operative antibiotics and aseptic techniques have become routine aspects of care in most major surgical procedures (4).

#### *Surgical site infections*

A surgical site infection (SSI) is “an infection related to an operative procedure that occurs at or near the surgical incision within 30 days of the procedure, or within 90 days if prosthetic material is implanted at surgery” (16). Surgical site infections are classified as follows (17);

#### Superficial incisional SSI

A superficial incisional SSI is an SSI which occurs within 30 days following an operative procedure (where day 1 = the procedure date) and which involves the skin and subcutaneous tissue of the incision. A patient with a superficial incisional SSI may have any of the following clinical features: pain, tenderness, swelling, erythema or drainage of pus occurring at the incision site and microbiological isolation of an organism from the incision (17). There are two specific types of superficial incisional SSIs (17):

**a. Superficial incisional primary (SIP):** This is a superficial incisional SSI that occurs in the primary incision in a patient that has had surgery involving one or more incisions such as a caesarian section incision (17).

**2. Superficial incisional secondary (SIS)** – this is a superficial incisional SSI that occurs in the secondary incision in a patient that has had surgery involving more than one incision such as the donor site incision for coronary bypass graft surgery (17).

#### Deep incisional SSI

A deep incisional SSI is an SSI which occurs within 30 or 90 days following the operative procedure

(where day 1 = the procedure date) and involves deep soft tissues of the incision such as fascial or muscle layers. A patient with a deep SSI may have any of the following clinical features: fever (>38°C), localized pain, tenderness, an abscess or purulent drainage at the site of deep incision; and microbiological isolation of an organism from the deep soft tissue of the incision (17). There are two specific types of deep incisional SSIs:

**1. Deep incisional primary (DIP):** This refers to a deep incisional SSI that occurs in a primary incision in a patient that has had surgery involving one or more incisions such as a caesarian section (17).

**2. Deep incisional secondary (DIS):** This is a deep incisional SSI that occurs in a secondary incision in a patient that has had surgery involving an operation with more than one incision for example the donor site incision for coronary bypass graft surgery (17).

#### *Organ space SSI*

This refers to an SSI that occurs within 30 or 90 days following surgery (where day 1 = the procedure date) and involves any part of the body deeper than the fascial/muscle layers that are opened or manipulated during the operative procedure (17). A patient with an organ space SSI may have any of the following clinical features: purulent drainage from a drain placed into the organ/space; organism(s) identified from fluid or tissue in the organ/space; an abscess or other evidence of infection involving the organ/space (17).

#### *Epidemiology of SSI*

SSIs are associated with patient morbidity and mortality and increased healthcare costs borne by the patient and healthcare providers (18, 19). In many low- and middle-income nations (LMICs), SSIs account for the highest frequency of healthcare-associated infections (20, 21). Globally the pooled incidence of SSI is estimated to be 2.5% [95% ci: 1.6, 3.7] (22). It ranges between 0.6 and 9.5% in Europe (23). In the USA a study showed an overall SSI rate of 1.9% (24). In the Asia-pacific region, the incidence of SSI varies per country with cumulative incidences of 2.8% in Australia, 2-9.7% in the Republic of Korea and China, 7.8% in south-east Asia and Singapore and 4% in China (20, 24, 25, 26, 27, 28). The pooled cumulative incidence of SSI in sub-Saharan Africa is 14.8% (29).

In Nigeria, a meta-analysis showed a cumulative incidence of SSI of 14.5%, [95% ci: 0.113–0.184] (30). SSIs in Nigeria were found to be quite common following colorectal surgeries (29.2%,

95% ci: 0.216–0.382), abdominal surgeries (20%, 95% ci: 0.064–0.649), and among the paediatric populations [29.6%, 95% ci: 0.136–0.529] (30). The north-eastern region of Nigeria has reported the highest rates of SSI (27.3%, 95% ci: 0.132–0.481) followed by the northcentral region [26.3%, 95% ci: 0.116–0.493] (30). However, the lowest rate of SSI was reported in the south-south region (8.0%, 95% ci: 0.065–0.098) of the country (30). The most frequently encountered type of SSI is the superficial incisional SSI, which occurred in 62.5% (ci: 0.333–0.848) of cases (30). The SSI incidence was also predominantly reported among patients with dirty wounds (52.7%, 95% ci: 0.367–0.682) and contaminated wounds [24.0%, 95% ci: 0.164–0.336] (30). Similarly, the highest incidence of 18.62% SSI was reported among patients aged 60 years and above followed by an incidence rate of 16.91% among patients aged <20 years (30).

#### *Risk factors for surgical site infections*

The risk factors for SSI include patient-related or endogenous factors (such as age, gender, weight, comorbidities etc.) and exogenous or process/procedural-related factors (such as duration of the surgery, adherence to infection control and prevention protocols etc.) (31).

Findings from a systematic review involving 57 studies from high-income countries and LMICs revealed that the following factors were associated with an increased risk of SSI: diabetes, prolonged duration of surgery, a high body mass index; a severe score according to the US National Nosocomial Infections Surveillance (NNIS) risk index; and a severe wound class (32). In a study conducted in Pakistan SSI was found to be more common in older patients (11.4% vs. 6.4%;  $p=0.009$ ); patients with more than 24 hours of preoperative hospital stay (11.2% vs. 6.4%;  $p=0.013$ ), in emergency surgeries (19.2% vs. 7.5%;  $p=0.0001$ ) and in procedures of longer duration ( $1.53 \pm 0.35$  vs  $2.57 \pm 0.17$ ;  $p<0.0001$ ) (33).

In a study done in Nigeria SSI was found to be more common in patients with diabetes mellitus (or = 1.2; ci: 0.159–9.109); elderly patients (or = 1.02; 95% ci: 0.993–1.055); prolonged duration of postoperative hospital stay (or = 1.07; 95% ci: 1.011–1.131) and cigarette smoking (or = 6.24; 95% ci: 0.274–142.15) (34). Another Nigerian study showed that age (>60 years), anaemia, obesity, number of individuals (> 6 individuals) in the operating room and duration of surgery were all significantly associated with the occurrence of SSIs (35).

#### *Microbiology of SSI*

A significant proportion of organisms responsible for SSIs are acquired endogenously or exogenously from the patient's environment (36, 37). In addition, the location of the health facility, duration of the surgery, site of the surgery and the presence (and level of adherence) of strict infection prevention and control protocols may also influence the type of pathogens involved in SSIs (38, 39).

However, bacteria are the organisms predominantly implicated in SSIs (38, 39). Among bacteria, *Staphylococcus aureus* is the most frequently isolated pathogen in SSIs (38, 39). However, in some studies gram-negative bacteria, such as *Escherichia Coli*, are the most frequently isolated pathogen (40).

Specifically, multi-drug resistant bacteria such as methicillin-resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase-negative *Staphylococci*, vancomycin-resistant enterococci and extended spectrum beta-lactamase (ESBL) producing Enterobacteriaceae are also very important causes of SSI particularly due to the significant morbidity and mortality associated with these organisms (41, 42).

#### *Preventing surgical site infections*

A variety of measures have been suggested for preventing surgical site infections. Examples include pre-operative bathing; decolonization with mupirocin ointment for the prevention of staphylococcus aureus infection in nasal carriers; screening of ESBI colonization; mechanical bowel preparation and the use of oral antibiotics; hair removal around the surgical site and surgical site preparation with antiseptic solutions (43). Other measures include the use of antimicrobial skin sealants; enhanced nutritional support; perioperative discontinuation of immunosuppressive agents; perioperative oxygenation, maintaining normal body temperature (normothermia); intensive perioperative blood glucose control in diabetics; maintenance of adequate circulating volume control/normovolemia; prophylactic negative pressure wound therapy; the use of antimicrobial -coated sutures; the use of laminar flow ventilation systems in the operating suite; and the administration of surgical antibiotic prophylaxis (43).

The World Health Organization has however adopted a multi-modal strategy to prevent the occurrence of SSIs (44). This strategy is a cohesive approach to ensure implementation and scalability of efforts to curb SSIs, particularly in resource-constrained settings which lack required

infrastructure such as laminar flow ventilation systems, water, sanitation and hygiene (wash) facilities etc (44). This strategy is made up of five components:

1. **System change:** adequate infrastructure, resources and protocols must be in place in efforts to curb SSI and also ensure the sustainability of these efforts (44). In addition, the right infrastructure, resources and protocols ensure that interventions to curb SSIs are streamlined, consistent and sustainable (44). For example, appropriate surgical techniques must go hand in hand with the availability of hand hygiene facilities in the theatre and post-operative wards (44).
2. **Training and education:** continuous training regarding infection prevention and control (IPC) protocols to curb SSIs is required for all healthcare staff who are involved in the management of surgical patients (44). Such training should present the rationale for IPC protocols in efforts to curb SSIs and incorporate behavioural change components to ensure improved outcomes in efforts to curb SSIs (44).
3. **Monitoring and feedback:** it is important to monitor and evaluate compliance with interventions to curb SSIs. For example compliance with hand hygiene, monitoring of blood glucose control in diabetics etc. In addition, monitoring and evaluation will also provide evidence on SSI rates and risk factors and use this knowledge to adjust or design interventions to curb SSIs (44).
4. **Reminders and communications for awareness raising:** for example, posters can be placed in strategic places in the hospital to remind or educate health workers patients and/or their relatives about surgical site infections (44).
5. **Institutional safety climate and culture change:** the leadership of each healthcare facility must create an enabling environment to curb SSIs. For example, acquiring the required infrastructure, ensuring compliance with IPC protocols and procuring antibiotics required for surgical prophylaxis (44).

However, a major component required to prevent surgical site infections is surgical antibiotic prophylaxis. This review further discusses surgical antibiotic prophylaxis (SAP).

#### *Principles of surgical prophylaxis*

The following are clearly defined principles regarding the practice of SAP.

#### **1. Indications for surgical prophylaxis**

To understand which surgeries, require SAP or not, surgical wounds, are classified according to their potential risk for infectious complications. This classification which has greatly facilitated the study and practice of surgical antibiotic prophylaxis is ranked in the following manner:

##### Classification of surgical wounds (1, 17)

**Class I/clean:** refers to uninfected operative wounds devoid of inflammation and in which the respiratory, alimentary, genital, or uninfected urinary tract is not entered. Examples include herniorrhaphy, mastectomy, cosmetic surgery, insertion of prosthesis (e.g. Hip replacement) or artificial devices (e.g. Heart valves) (1, 17).

**Class II/ clean-contaminated:** this is an operative wound in which the respiratory, alimentary, genital, or urinary tracts are entered under controlled conditions and without unusual contamination. Examples include laryngectomy, uncomplicated appendectomy, cholecystectomy, and transurethral resection of the prostate gland (1, 17).

**Class III/ contaminated:** these include open, fresh, accidental wounds, operations with major breaks in sterile technique (e.g., open cardiac massage), or gross spillage from the gastrointestinal tract and incisions in which acute, non-purulent inflammation is encountered are included in this category (1, 17). Examples include large bowel resection, and biliary or genitourinary tract surgery with infected bile or urine (1, 17).

**Class IV/dirty-infected wounds:** these are old traumatic wounds with retained devitalized tissue and those that involve existing clinical infection or perforated viscera. This definition suggests that the organisms causing postoperative infection were present in the operative field before the operation (17).

Based on the classification above SAP is uniformly recommended for all clean-contaminated and contaminated procedures and operations involving the insertion of an artificial device or prosthetic material (1).

#### **2. Choice of antibiotics for surgical prophylaxis**

The following factors are considered in selecting antibiotics for SAP. These include: ensuring that the choice of antibiotics provides coverage against the expected endogenous flora at the surgical site; antibiotic penetration into the surgical site; antimicrobial costs to the patient, the existence of

patient allergies to any of the antibiotics; a knowledge of the ecology of local nosocomial wound pathogens, and ensuring appropriate antibiotic dosing and administration (4). In addition, where two or more antibiotics (which are equal in the antibacterial spectrum, efficacy, toxicity, and ease of administration) are available the least expensive drug should be chosen (1, 5).

### 3. Route of antibiotic administration

The intravenous route is the usual mode of administering prophylactic antibiotics. They are usually administered as a bolus at induction of anaesthesia to ensure adequate tissue concentrations at the time of surgical incision (1). The intramuscular route of administration is a less commonly used route than the intravenous route. Intramuscularly administered prophylactic antibiotics are given at the time of pre-medication so that peak tissue levels are attained at the most critical time (i.e. The time of surgical incision) (1). Other routes of administration include oral or rectal routes which are commonly used in bowel surgeries. Topical antibiotics are not recommended routes for surgical prophylaxis, with the exceptions of ophthalmic or burns surgery (1). The use of antibiotic-impregnated cement placed directly into the operative wound (as a local antimicrobial *brachytherapy*) has also been deployed as a method of SAP particularly in procedures involving the replacement of infected prosthetic joints (45, 46). It has been significantly associated with a reduction in surgical site infections (45, 46).

### 4. Timing of antibiotic administration

Appropriately timed antibiotic prophylaxis is defined as “the delivery of the antibiotic within one hour before incision, with the exception that vancomycin and the fluoroquinolones should be given within two hours before incision because of the need for a longer infusion time” (4, 8). Inappropriate timing of antibiotic prophylaxis has been associated with suboptimal tissue levels and an increased risk for surgical site infections (4). For example, Classen and colleagues noted that the risk of SSI was reduced when antibiotics were administered within two hours before incision (47). Also, investigators involved in the trial to reduce antimicrobial prophylaxis errors (trape) examined the association between SSI and timing of prophylaxis in cardiac, orthopaedic, and hysterectomy patients and found that SSI risk was lowest in those patients who received prophylaxis within 30 minutes (if given cephalosporins) or within 1 hour (if given vancomycin or a

fluoroquinolone) before incision. They also found that post-incision administration of prophylaxis was associated with a significantly increased risk for SSI (48).

However, oral or rectal antibiotics are given earlier than the time frames mentioned above, to achieve optimal tissue concentrations (1). For example, metronidazole suppositories, which are commonly used in bowel surgery, must be given two to four hours before surgery (1).

### 5. Duration of antibiotic administration

The discontinuation of prophylactic antibiotics within 24 hours of completing surgery is recommended by most guidelines for SAP (8). For surgeries not exceeding four hours a single dose of the antibiotic is usually sufficient while in prolonged surgeries exceeding four hours, further antibiotic doses may be required to maintain the concentration, particularly if the antibiotic has a short half-life (1, 8).

For cardiac surgery, experts recommend continuing prophylaxis for 48 hours, based on concerns that more data are needed before uniformly recommending a shorter duration of antibiotic administration (49). The prolonged administration of antibiotic prophylaxis in the postoperative period (e.g., until surgical drains have been removed) does not improve efficacy and increases toxicity, cost and drug resistance (4). For example, Harbath and co-workers found that prolonged antibiotic prophylaxis (>48 hours post-incision) was significantly associated with an increased risk of acquiring an antibiotic-resistant pathogen (50). Another study conducted by Arrigan et al (2007) in Zambia found that prolonged administration of surgical prophylaxis beyond 24 hours resulted in no benefit to the patients and was associated with an increased risk of surgical site infections, increased length of hospitalization and increased costs borne by the patient (51).

#### *Challenges, regarding the practice of surgical antibiotic prophylaxis*

Globally the use of surgical prophylaxis is supported in several surgical procedures e.g., gastrointestinal, oropharyngeal, vascular, orthopaedic, etc (52). However, the practice of SAP is fraught with variations and controversies regarding issues such as indications for SAP, choice of antibiotics, timing, and duration of prophylaxis (3, 4, 53). These variations and controversies are essentially due to several factors:

1. **First** is a lack of institutional or national guidelines for SAP. This problem is particularly

pronounced in low- and middle-income countries (Imics) with weak governance structures and fragile health systems (54, 55).

2. **Second** is the poor compliance by surgeons with available guidelines and recommendations on SAP. The problem of compliance with guidelines is global (8, 53), as evidenced by previous studies in several institutions which have shown wide ranges of compliance ranging from 0% to 71.9% with the majority of studies showing compliance rates below 50% (56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68). For example, findings from a multi-centre audit of elective procedures in thirteen hospitals in the Netherlands showed that only 28% of the procedures achieved full adherence to all parameters of the local hospital guidelines (i.e. Choice of antibiotic, duration, dose, dosing interval and timing of first dose (54).also, a study done in a hospital in Doha, Qatar showed a compliance rate of 68.5% and 40.7% regarding the hospital's guidelines on antibiotic selection and duration of SAP respectively (69). Another study in the united arab emirates showed a compliance rate of 30.4% regarding international guidelines on dosage (66) while a study done in Sudan showed an overall compliance rate of 2.7% with international guidelines (70). In Nigeria, previous studies have shown compliance rates less than 50% (71, 72, 73, 74).

A lack of guidelines or poor compliance with available guidelines contributes to the inappropriate use of antibiotics for SAP in several healthcare facilities, as a variety (rather than a uniform regime) of prescriptions for SAP will emanate from each surgical unit (49). Sadly, inappropriate antibiotic use exerts adverse clinical and economic impacts on surgical patients (75, 76). While the adverse clinical impacts include increased hospital costs borne by the patients, the adverse clinical impacts include the occurrence of surgical site infections, prolonged hospital stay, increased consumption of antibiotics, the emergence of antibiotic resistance, morbidity and mortality (75, 76). Also, previous studies have shown that in the absence of guidelines the prescribing patterns of physicians for SAP tend to be based on diverse rationales such as personal preferences, medical literature (77), discussions with colleagues (78) etc. In addition, there is a misconception among some surgeons that prolonged regimens with multiple antibiotics are more effective than short courses of narrow-

spectrum antibiotics in reducing or eliminating the risk of surgical site infections (79).

Improving the practice of SAP will involve developing institutional guidelines. The approach to developing these guidelines should be multi-disciplinary and must involve the hospital's antimicrobial stewardship team, surgeons, microbiologists, anaesthetists, and pharmacists (8, 52).

To ensure compliance with guidelines it will be necessary to conduct regular institutional audits on the practice of SAP (5). An audit is defined as "a quality improvement process that seeks to improve patient care and outcomes through a systematic review of all aspects of care against explicit criteria and the implementation of change. Where indicated, changes are implemented at an individual, team, or service level and further monitoring is used to confirm improvement in healthcare delivery" (80).

Continuing medical education (e.g., seminars, courses, workshops etc.) Will also be necessary to ensure updates and exposure to current best practices regarding surgical antibiotic prophylaxis (81, 82).

More cost-intensive measures may include the deployment of information technology (e.g., computer-based order systems) to avoid excessive duration of surgical antibiotic prophylaxis (83).

The commitment of a hospital's leadership is integral to improving the practice of surgical antibiotic prophylaxis (84). Examples of leadership commitment include making available the required human, financial, and information technology resources (84).

Improving the practice of surgical antibiotic prophylaxis is however not a standalone effort but should go alongside other quality improvement initiatives (i.e., antimicrobial stewardship, infection prevention and control, improved diagnostic microbiology capacity, patient safety etc.) In the hospital (81).

### Conclusion

Surgical antibiotic prophylaxis (SAP) is useful in preventing surgical site infections and improving post-operative outcomes (1). However, the practice of SAP is challenged by poor guideline adherence resulting in adverse clinical and economic outcomes for surgical patients (3, 4, 53, 75, 76). Improving the practice of SAP will include developing guidelines, conducting regular audits to assess guidelines and continuing medical education (5, 8, 52, 80, 81, 82). The leadership of a healthcare facility must also be committed to

improving the practice of SAP by making available the needed human, financial, and material resources (84). Finally, efforts to improve the practice of SAP in a healthcare facility must be deployed in consonance with other quality improvement initiatives such as antimicrobial stewardship, infection prevention and control, and improved diagnostic microbiology capacity (81).

#### List of Abbreviations

DIP: Deep Incisional Primary  
DIS: Deep Incisional Secondary  
IPC: Infection Prevention and Control  
LMIC: Low- and Middle-Income Countries  
SAP: Surgical Antibiotic Prophylaxis  
SIP: Superficial Incisional Primary  
SIS: Superficial Incisional Secondary  
SSI: Surgical Site Infection  
Trape: Trial to reduce antimicrobial prophylaxis errors  
WASH: Water, Sanitation and Hygiene

#### Declarations

*Ethics approval and consent to participate*  
Not applicable.

#### *Consent for publication*

The authors hereby transfer all copyright ownership exclusively to the journal, if the journal publishes this work. The authors give consent for the publication of the work under the Creative Commons Attribution-Non-Commercial 4.0 license.

#### *Availability of data and materials*

The articles used in this study are publicly available.

#### *Competing interests*

The authors have declared no conflict of interest.

#### *Funding*

The authors received no research funding.

#### *Authors' contributions*

IIO conceptualized, designed the study, and drafted the manuscript.

#### Acknowledgement

None.

#### Reference

1. Munckhof W. Antibiotics for surgical prophylaxis AustPrescr 2005; 28:38–40. <https://doi.org/10.18773/austprescr.2005.030>

2. Khan A K A, P V M, Rashed MR, Banu G. A Study on the Usage Pattern of Antimicrobial Agents for the Prevention of Surgical Site Infections (SSIs) in a Tertiary Care Teaching Hospital. J Clin Diagn Res. 2013;7(4):671-4. <https://doi.org/10.7860/JCDR/2013/5323.2878>
3. Finland M. Antibacterial agents: uses and abuses in treatment and prophylaxis. R I Med J. 1960;43:499-504.
4. Talbot TR. Surgical Site Infections and Antimicrobial Prophylaxis. In: Mandell, Douglas and Bennett's Principles and Practice of Infectious Diseases. 8th ed. Philadelphia: Churchill Livingstone; 2015. p. 4684-4698. Vol 2
5. Sinclair A, Bridgewater, B. How to Set Up a Prospective Surgical Audit In Wilcox DT, Godbole PP and Koyle MA, editors. Paediatric Urology: Surgical Complications and Management. Chichester: Blackwell Publishing; 2008. p.1-7
6. van Kasteren ME, Mannien J, Kullberg BJ, de Boer AS, Nagelkerke NJ, Ridderhof M, Wille JC, Gyssens IC. Quality improvement of surgical prophylaxis in Dutch hospitals: evaluation of a multi-site intervention by time series analysis. Journal of Antimicrobial Chemotherapy. 2005 Dec 1;56(6):1094-102. <https://doi.org/10.1093/jac/dki374>
7. Fukatsu K, Saito H, Matsuda T, Ikeda S, Furukawa S, Muto T. Influences of type and duration of antimicrobial prophylaxis on an outbreak of methicillin-resistant Staphylococcus aureus and on the incidence of wound infection. Arch Surg. 1997 Dec;132(12):1320-5. <https://doi.org/10.1001/archsurg.1997.01430360066012>
8. Bratzler DW, Hunt DR. The surgical infection prevention and surgical care improvement projects: national initiatives to improve outcomes for patients having surgery. Clin Infect Dis. 2006;43(3):322-30. <https://doi.org/10.1086/505220>
9. Steinberg JP, Braun BI, Hellinger WC, Kusek L, Bozikis MR, Bush AJ, Dellinger EP, Burke JP, Simmons B, Kritchevsky SB; Trial to Reduce Antimicrobial Prophylaxis Errors (TRAPE) Study Group. Timing of antimicrobial prophylaxis and the risk of surgical site infections: results from the Trial to Reduce Antimicrobial Prophylaxis Errors. Ann Surg. 2009;250(1):10-6. <https://doi.org/10.1097/SLA.0b013e3181ad5fca>

10. Culbertson WR, Altemeier WA, Gonzalez LL, Hill EO. Studies on the epidemiology of postoperative infection of clean operative wounds. *Ann Surg.* 1961 Oct;154(4):599-610.
11. Howe CW, Marston AT. A study on sources of postoperative staphylococcal infection. *Surg Gynecol Obstet.* 1962;115: 266-75. <https://doi.org/10.1097/00000658-196110000-00010>
12. Burke JF. Identification of the sources of Staphylococci contaminating the surgical wound during operation. *Ann Surg.* 1963;158(5):898-904. <https://doi.org/10.1097/00000658-196311000-00021>
13. Howes EL. Prevention of wound infection by the injection of nontoxic antibacterial substances. *Ann Surg.* 1946;124 (2):268-276.
14. Miles AA, Miles EM, Burke J. The value and duration of defence reactions of the skin to the primary lodgement of bacteria. *Br J Exp Pathol.* 1957;38(1):79-96.
15. Burke JF. The effective period of preventive antibiotic action in experimental incisions and dermal lesions. *Surgery.* 1961; 50:161-8.
16. Agency for Healthcare Research and Quality. Patient Safety Network. Surgical Site Infections. [https://psnet.ahrq.gov/primer/surgical-site-infections#:~:text=Surgical%20site%20infection%20\(SSI\)%E2%80%94,among%20the%20most%20common%20preventable](https://psnet.ahrq.gov/primer/surgical-site-infections#:~:text=Surgical%20site%20infection%20(SSI)%E2%80%94,among%20the%20most%20common%20preventable)
17. United States Centers for Disease Control. National Healthcare Safety Network. Surgical Site Infection Event. 2023. [https://www.cdc.gov/nhsn/PDFs/pscManual/9\\_pscSSICurrent.pdf?agree=yes&next=Accept](https://www.cdc.gov/nhsn/PDFs/pscManual/9_pscSSICurrent.pdf?agree=yes&next=Accept)
18. Hou Y, Collinsworth A, Hasa F, Griffin L. Incidence and impact of surgical site infections on length of stay and cost of care for patients undergoing open procedures. *Surgery Open Science.* 2023 Jan 1;11:1-8. <https://doi.org/10.1016/j.sopen.2022.10.004>
19. Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA, Crosby C. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *Journal of Hospital Infection.* 2017 May 1;96(1):1-5. <https://doi.org/10.1016/j.jhin.2017.03.004>
20. World Health Organization. Report on the burden of endemic health care-associated infection worldwide. A systematic review of the literature. Geneva: World Health Organization. 2011 [https://iris.who.int/bitstream/handle/10665/80135/9789241501507\\_eng.pdf?sequence=1](https://iris.who.int/bitstream/handle/10665/80135/9789241501507_eng.pdf?sequence=1)
21. Allegranzi B, Nejad SB, Combescure C, Graafmans W, Attar H, Donaldson L, Pittet D. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *The Lancet.* 2011 Jan 15;377(9761):228-41. [https://doi.org/10.1016/S0140-6736\(10\)61458-4](https://doi.org/10.1016/S0140-6736(10)61458-4)
22. Mengistu DA, Alemu A, Abdulkadir AA. Global Incidence of Surgical Site Infection Among Patients: Systematic Review and Meta-Analysis. *Inquiry.* 2023; 60:469580231162549. <https://doi.org/10.1177/00469580231162549>
23. European Centres for Disease Control. Annual Epidemiological Meeting for 2018 to 2020. <https://www.ecdc.europa.eu/en/publications-data/monitoring/all-annual-epidemiological-reports>
24. Mu Y, Edwards JR, Horan TC, Berrios-Torres SI, Fridkin SK. Improving risk-adjusted measures of surgical site infection for the National Healthcare Safety Network. *Infection Control & Hospital Epidemiology.* 2011 Oct;32(10):970-86. <https://doi.org/10.1086/662016>
25. Takesue Y, Watanabe A, Kusachi S, Matsumoto T, Iwamoto A, Totsuka K, Sunakawa K, Yagisawa M, Sato J, Oguri T, Nakanishi K. Nationwide surveillance of antimicrobial susceptibility patterns of pathogens isolated from surgical site infections (SSI) in Japan. *Journal of Infection and Chemotherapy.* 2012 Jan 1;18(6):816-26. <https://doi.org/10.1007/s10156-012-0509-1>
26. Curcio D, Cane A, Fernández F, Correa J. Surgical site infection in elective clean and clean-contaminated surgeries in developing countries. *Int J Infect Dis.* 2019;80:34–45. <https://doi.org/10.1016/j.ijid.2018.12.013>
27. Worth LJ, Bull AL, Spelman T, Brett J, Richards MJ. Diminishing surgical site infections in Australia: time trends in infection rates, pathogens and antimicrobial resistance using a comprehensive Victorian surveillance program, 2002–2013. *infection control & hospital epidemiology.* 2015 Apr;36(4):409-16. <https://doi.org/10.1017/ice.2014.70>
28. Ling ML, Apisarnthanarak A, Madriaga G. The burden of healthcare-associated infections in Southeast Asia: a systematic literature review and meta-analysis. *Clin Infect Dis.* 2015;60(11):1690–9. <https://doi.org/10.1093/cid/civ095>

29. Ngah JE, Bénet T, Djibrilla Y. Incidence of surgical site infections in sub-Saharan Africa: systematic review and meta-analysis. *The Pan African Medical Journal*. 2016 Jun 29;24:171-54. <https://doi.org/10.11604/pamj.2016.24.171.9754>
30. Olowo-Okere A, Ibrahim YK, Olayinka BO, Ehinmidu JO. Epidemiology of surgical site infections in Nigeria: a systematic review and meta-analysis. *Nigerian Postgraduate Medical Journal*. 2019 Jul 1;26(3):143-51. [https://doi.org/10.4103/npmj.npmj\\_72\\_19](https://doi.org/10.4103/npmj.npmj_72_19)
31. Cheadle WG. Risk factors for surgical site infection. *Surg Infect (Larchmt)*. 2006;7 (Suppl 1): S7-11. <https://doi.org/10.1089/sur.2006.7.s1-7>
32. Korol E, Johnston K, Waser N, Sifakis F, Jafri HS, Lo M, Kyaw MH. A systematic review of risk factors associated with surgical site infections among surgical patients. *PloS one*. 2013 Dec 18;8(12):e83743. <https://doi.org/10.1371/journal.pone.0083743>
33. Ansari S, Hassan M, Barry HD, Bhatti TA, Hussain SZ, Jabeen S, Fareed S. Risk factors associated with surgical site infections: a retrospective report from a developing country. *Cureus*. 2019 Jun 2;11(6). <https://doi.org/10.7759/cureus.4801>
34. Olowo-Okere A, Ibrahim YKE, Sani AS, Olayinka BO. Occurrence of Surgical Site Infections at a Tertiary Healthcare Facility in Abuja, Nigeria: A Prospective Observational Study. *Med Sci (Basel)*. 201;6(3):60. <https://doi.org/10.3390/medsci6030060>
35. Nwankwo, E., Ibeh, I., & Enabulele, O. Incidence and risk factors of surgical site infection in a tertiary health institution in Kano, Northwestern Nigeria. *International Journal of Infection Control*. 2012; 8:4. <https://doi.org/10.3396/ijic.v8i4.10573>
36. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control*.
37. Cooper RA. Surgical site infections: epidemiology and microbiological aspects in trauma and orthopaedic surgery. *Int Wound J* 2013; 10 (suppl. 1):3–8. <https://doi.org/10.1111/iwj.12179>
38. National Nosocomial Infections Surveillance System. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am J Infect Control* 2004;32:470–85. <https://doi.org/10.1016/S0196655304005425>
39. Misteli H, Widmer AF, Rosenthal R, Oertli D, Marti WR, Weber WP. Spectrum of pathogens in surgical site infections at a Swiss university hospital. *Swiss Med Wkly* 2011;140:w13146. <https://doi.org/10.4414/smww.2011.13146>
40. Okoro KA, Ede O, Iyidobi EC, Enweani UU, Nwadinigwe CU, Eyichukwu GO, Anyaehie UE, Ahaotu FN, Ezech RC. The Bacteriological Profile of Surgical Site Infections in Orthopaedic Implant Surgeries in South-East Nigeria. *Journal of Biosciences and Medicines*. 2019 Aug 30;7(09):19.
41. Krulerink M, Kievit J, Marang-van de Mheen PJ. Evaluation of routinely reported surgical site infections against microbiological culture results: a tool to identify patient groups where diagnosis and treatment may be improved. *BMC Infect Dis* 2009;9:176doi: <https://doi.org/10.1186/1471-2334-9-176>
42. Martínez-Pastor JC, Vilchez F, Pitart C, Sierra JM, Soriano A. Antibiotic resistance in orthopaedic surgery: acute knee prosthetic joint infections due to extended-spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae. *European journal of clinical microbiology & infectious diseases*. 2010 Aug;29:1039-41. <https://doi.org/10.1007/s10096-010-0950-y>
43. World Health Organization. Global guidelines for the prevention of surgical site infection, second edition. Geneva: World Health Organization; 2018. <https://www.ncbi.nlm.nih.gov/books/NBK536404>
44. World Health Organization. Preventing surgical site infections: implementation approaches for evidence-based recommendations. Geneva: World Health Organization; 2018. <https://iris.who.int/bitstream/handle/10665/273154/9789241514385-eng.pdf?sequence=1>
45. Winger DA, Fass RJ. Antibiotic-impregnated cement and beads for orthopedic infections. *Antimicrob Agents Chemother*. 1996;40(12):2675-9. <https://doi.org/10.1128/AAC.40.12.2675>
46. Goodell JA, Flick AB, Hebert JC, Howe JG. Preparation and release characteristics of tobramycin-impregnated polymethylmethacrylate beads. *Am J Hosp Pharm*. 1986;43(6):1454-61.
47. Classen DC, Evans RS, Pestotnik SL, Horn SD, Menlove RL, Burke JP. The timing of prophylactic administration of antibiotics and the risk of surgical-wound infection. *N Engl J*

- Med. 1992;326(5):281-6.  
<https://doi.org/10.1056/NEJM199201303260501>
48. Steinberg JP, Braun BI, Hellinger WC, Kusek L, Bozikis MR, Bush AJ, Dellinger EP, Burke JP, Simmons B, Kritchevsky SB; Trial to Reduce Antimicrobial Prophylaxis Errors (TRAPE) Study Group. Timing of antimicrobial prophylaxis and the risk of surgical site infections: results from the Trial to Reduce Antimicrobial Prophylaxis Errors. *Ann Surg.* 2009 Jul;250(1):10-6.  
<https://doi.org/10.1097/SLA.0b013e3181ad5fca>
  49. Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. *Ann Thorac Surg.* 2006 ;81(1):397-404.  
<https://doi.org/10.1016/j.athoracsur.2005.06.034>
  50. Harbarth S, Samore MH, Lichtenberg D, Carmeli Y. Prolonged antibiotic prophylaxis after cardiovascular surgery and its effect on surgical site infections and antimicrobial resistance. *Circulation.* 2000;101(25):2916-21.  
<https://doi.org/10.1161/01.cir.101.25.2916>
  51. Arrigan M, Halley B, Hughes P, Mcmenamin L, O'Sullivan K. Pharmacoeconomic Analysis of Peri-Surgical Antibiotics and Surgical Site Infections in Livingstone General Hospital, Zambia TSMJ, 2007; 8 (1): 71-75.  
<https://ojs.tchpc.tcd.ie/index.php/tsmj/article/view/1835>
  52. World Health Organization. WHO guidelines for safe surgery 2009: safe surgery saves lives.  
[https://apps.who.int/iris/bitstream/handle/10665/44185/9789241598552\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/44185/9789241598552_eng.pdf)
  53. Ng RS, Chong CP. Surgeons' adherence to guidelines for surgical antimicrobial prophylaxis - a review. *Australas Med J.* 2012;5(10):534-40.  
<https://doi.org/10.4066/AMJ.2012.1312>
  54. Nnadozie UU, Umeokonkwo CD, Maduba CC, Igwe-Okomiso D, Onah CK, Madubueze UC, Anikwe CC, Versporten A, Pauwels I, Goossens H, Ogbuanya AU, Oduyebo OO, Onwe EO. Antibiotic use among surgical inpatients at a tertiary health facility: a case for a standardized protocol for presumptive antimicrobial therapy in the developing world. *Infect Prev Pract.* 2020;2(4):100078.  
<https://doi.org/10.1016/j.infpip.2020.100078>
  55. Siachalinga L, Godman B, Mwita JC, Sefah IA, Ogunleye OO, Masele A, Lee IH. Current Antibiotic Use Among Hospitals in the sub-Saharan Africa Region; Findings and Implications. *Infect Drug Resist.* 2023;16:2179-2190.  
<https://doi.org/10.2147/IDR.S398223>
  56. Al-Momany NH, Al-Bakri AG, Makahleh ZM, Wazaify MM. Adherence to international antimicrobial prophylaxis guidelines in cardiac surgery: a Jordanian study demonstrates need for quality improvement. *J Manag Care Pharm.* 2009;15(3):262-71.  
<https://doi.org/10.18553/jmcp.2009.15.3.262>
  57. Lallemand S, Thouverez M, Bailly P, Bertrand X, Talon D. Non-observance of guidelines for surgical antimicrobial prophylaxis and surgical-site infections. *Pharm World Sci.* 2002;24(3):95-9.  
<https://doi.org/10.1023/a:1016122202439>
  58. Ozgun H, Ertugrul BM, Soyder A, Ozturk B, Aydemir M. Peri-operative antibiotic prophylaxis: adherence to guidelines and effects of educational intervention. *Int J Surg.* 2010;8(2):159-63.  
<https://doi.org/10.1016/j.ijsu.2009.12.005>
  59. Gorecki P, Schein M, Rucinski JC, Wise L. Antibiotic administration in patients undergoing common surgical procedures in a community teaching hospital: the chaos continues. *World J Surg.* 1999;23(5):429-32.  
<https://doi.org/10.1007/pl00012319>
  60. Gul YA, Hong LC, Prasanna S. Appropriate antibiotic administration in elective surgical procedures: still missing the message. *Asian J Surg.* 2005;28(2):104-8.  
[https://doi.org/10.1016/S1015-9584\(09\)60272-4](https://doi.org/10.1016/S1015-9584(09)60272-4)
  61. Bratzler DW, Houck PM, Richards C, Steele L, Dellinger EP, Fry DE, Wright C, Ma A, Carr K, Red L. Use of antimicrobial prophylaxis for major surgery: baseline results from the National Surgical Infection Prevention Project. *Arch Surg.* 2005;140(2):174-82.  
<https://doi.org/10.1001/archsurg.140.2.174>
  62. Bailly P, Lallemand S, Thouverez M, Talon D. Multicentre study on the appropriateness of surgical antibiotic prophylaxis. *J Hosp Infect.* 2001;49(2):135-8.  
<https://doi.org/10.1053/jhin.2001.1064>
  63. Tourmousoglou CE, Yiannakopoulou ECh, Kalapothaki V, Bramis J, St Papadopoulos J. Adherence to guidelines for antibiotic prophylaxis in general surgery: a critical appraisal. *J Antimicrob Chemother.* 2008;61(1):214-8.  
<https://doi.org/10.1093/jac/dkm406>

64. Askarian M, Moravveji AR, Mirkhani H, Namazi S, Weed H. Adherence to American Society of Health-System Pharmacists surgical antibiotic prophylaxis guidelines in Iran. *Infect Control Hosp Epidemiol.* 2006;27(8):876-8. <https://doi.org/10.1086/506405>
65. Heineck I, Ferreira MB, Schenkel EP. Prescribing practice for antibiotic prophylaxis for cesarean section in a teaching hospital in Brazil. *Am J Infect Control.* 2002;30(6):341-5. <https://doi.org/10.1067/mic.2002.123347>
66. Hosoglu S, Sunbul M, Erol S, Altindis M, Caylan R, Demirdag K, Ucmak H, Mendes H, Geyik MF, Turgut H, Gundes S. A national survey of surgical antibiotic prophylaxis in Turkey. *Infection Control & Hospital Epidemiology.* 2003 Oct;24(10):758-61. <https://doi.org/10.1086/502127>
67. Matti PR, Querol RC, Antonio-Velmonte M, de Vera R, Alejandria M. Prescribing practices of surgeons and factors that limit adherence to the Philippine College of Surgeons Clinical Practice Guidelines on Antimicrobial Prophylaxis for elective surgical procedures at the UP-PGH surgical wards. *Phil J Microbiol Infect Dis.* 2002;31(3):107-24.
68. Abdel-Aziz A, El-Menyar A, Al-Thani H, Zarour A, Parchani A, Asim M, El-Enany R, Al-Tamimi H, Latifi R. Adherence of surgeons to antimicrobial prophylaxis guidelines in a tertiary general hospital in a rapidly developing country. *Advances in Pharmacological and Pharmaceutical Sciences.* 2013 Jan 1;2013. <https://doi.org/10.1155/2013/842593>
69. Abu-Gharbieh E, Fahm S Adherence to Surgical Site Infection Guidelines in Cardiac Surgery in a Tertiary Hospital in Dubai, United Arab Emirates *Tropical Journal of Pharmaceutical Research* August 2012; 11 (4): 657-664.
70. Ajibade A, Akinniyi OT. Timing of prophylactic antibiotic administration in an orthopaedic hospital in a developing country *Sahel Medical Journal* 2013; 16(4):144-147. <https://doi.org/10.4103/1118-8561.125555>
71. Abubakar U. PSU41 Compliance With Surgical Antibiotic Prophylaxis For Obstetric And Gynecologic Surgeries In Nigeria: A Retrospective Study. *Value in Health* 2019;22(3): S899. <https://doi.org/10.1016/j.jval.2019.09.2625>
72. Fatiregun AA, Lochlainn LN, Kaboré L, Dosumu M, Isere E, Olaoye I. Missed opportunities for vaccination among children aged 0–23 months visiting health facilities in a southwest State of Nigeria. 2019. *Plos one.* 2021;16(8):e0252798 <https://doi.org/10.1371/journal.pone.0252798>
73. Oshikoya KA, Ogunyinka IA, Adamaigbo C, Olowo-Okere A. Surgical antimicrobial prophylaxis and its dose appropriateness among paediatric patients in a Nigerian teaching hospital. *J Chemother.* 2019;31(6):329-342. <https://doi.org/10.1080/1120009X.2019.1615725>
74. Wood C, Kostkova P, Olufemi O, Soriano D, Ogunsola F, Lefevre-Lewis C, Kpokiri E, Shallcross L. Understanding non-compliance with surgical antibiotic prophylaxis prescribing guidance, in Nigeria. *European Journal of Public Health.* 2019; 29(4):ckz186.088, <https://doi.org/10.1093/eurpub/ckz186.088>
75. Fraser A, Paul M, Almanasreh N, Tacconelli E, Frank U, Cauda R, Borok S, Cohen M, Andreassen S, Nielsen AD, Leibovici L. Benefit of appropriate empirical antibiotic treatment: thirty-day mortality and duration of hospital stay. *The American journal of medicine.* 2006 Nov 1;119(11):970-6. <https://doi.org/10.1016/j.amjmed.2006.03.034>
76. Janssen J, Afari-Asiedu S, Monnier A, Abdulai MA, Tawiah T, Wertheim H, Baltussen R, Asante KP. Exploring the economic impact of inappropriate antibiotic use: the case of upper respiratory tract infections in Ghana. *Antimicrobial Resistance & Infection Control.* 2022 Apr 1;11(1):53. <https://doi.org/10.1186/s13756-022-01096-w>
77. Gul YA, Lian LH, Jabar FM, Moissinac K. Antibiotic prophylaxis in elective colorectal surgery. *ANZ J Surg.* 2002;72(10):275-8. <https://doi.org/10.1046/j.1445-2197.2002.02365.x>
78. Lerano C, Thursky K, Peel T, Rajkhowa A, Marshall C, Ayton D. Influences on surgical antimicrobial prophylaxis decision making by surgical craft groups, anaesthetists, pharmacists and nurses in public and private hospitals. *PLoS One.* 2019; 14(11):e0225011. <https://doi.org/10.1371/journal.pone.0225011>
79. Parulekar L , Soman R, Singhal T, Rodrigues C, Dastur FD and Mehta AJ. How good is compliance with surgical antibiotic prophylaxis guidelines in a tertiary care private hospital in India? A prospective study. *Indian J Surg* 2009. 71(1):15–18. <https://doi.org/10.1007/s12262-009-0004-9>

80. National Institute for Clinical Excellence. Principles for best practice in clinical audit. Oxford: Radcliffe Medical Press; 2002.
81. Haney V, Maman S, Prozesky J, Bezinover D, Karamchandani K. Improving intraoperative administration of surgical antimicrobial prophylaxis: a quality improvement report. *BMJ Open Qual.* 2020;9(3):e001042. <https://doi.org/10.1136/bmjog-2020-001042>
82. McHugh SM, Corrigan MA, Dimitrov BD, Cowman S, Tierney S, Hill AD, Humphreys H. Preventing infection in general surgery: improvements through education of surgeons by surgeons. *Journal of Hospital Infection.* 2011 Aug 1;78(4):312-6. <https://doi.org/10.1016/j.jhin.2011.03.023>
83. Haynes K, Linkin DR, Fishman NO, Bilker WB, Strom BL, Pifer EA, Hennessy S. Effectiveness of an information technology intervention to improve prophylactic antibacterial use in the postoperative period. *J Am Med Inform Assoc.* 2011;18(2):164-8. <https://doi.org/10.1136/jamia.2009.002998>
84. CDC. Core Elements of Hospital Antibiotic Stewardship Programs. Atlanta, GA: US Department of Health and Human Services, CDC. 2019. <https://www.cdc.gov/antibiotic-use/core-elements/hospital.html>